


AMENDMENTS TO THE CLAIMS


This Listing of Claims will replace all prior versions, and listings of claims in the application:

LISTING OF CLAIMS:


1. (Currently amended) A process for producing of a silicone compound represented by the following formulas (a) and/or (a'),



comprising reacting a carboxylic acid represented by the following formula (a2)




to an epoxy silane represented by the following formula (a1)



obtaining a metal salt of the carboxylic acid represented by the general formula (a2), and
~~subsequently mixing the epoxy silane represented by the general formula (a1), the~~
~~carboxylic acid represented by formula (a2), the metal salt of the carboxylic acid represented by~~
~~the general formula (a2) and water;~~


wherein the reaction is carried out in presence of 0.050.3 wt% or more water in said reaction system, wherein A denotes a siloxanyl group, R¹ denotes a substituent with 1 to 20 carbons having

{W:21089\0207151-us0\01628859.DOC  }

a polymerizable group, R^2 to R^4 respectively and independently denote hydrogen, a substituted or unsubstituted substituent with 1 to 20 carbons, or $-X-A$, and X denotes a substituted or unsubstituted divalent substituent with 1 to 20 carbons.

2. (Previously presented) A process for producing of a silicone compound, wherein the silicone compound obtained according to Claim 1 is purified by a silica gel column or an alumina column.


3. (Withdrawn) A silicone compound obtained by the process according to Claim 1, wherein the siloxanyl group A is an atomic group represented by the following formula (b),




wherein, A^1 to A^{11} respectively and independently denote any one of hydrogen, a substituted or unsubstituted alkyl group with 1 to 20 carbon atoms and a substituted or unsubstituted aryl group with 6 to 20 carbons, n denotes an integer of 0 to 200, a, b and c denote respectively and independently an integer of 0 to 20, and $n = a = b = c = 0$ is not included.

4. (Withdrawn) A silicone compound according to Claim 3, wherein the siloxanyl group A is selected from the group consisting of tris(trimethylsiloxy)silyl group, bis(trimethylsiloxy)methylsilyl group and trimethylsiloxydimethylsilyl group.

5. (Withdrawn) A silicone compound in which a content of a compound represented by the following general formula (y) is 0.4% or more and 3% or less,




and the purity of the silicone compound represented by the following general formulas (a) and/or (a') is 87% or more,




wherein A denotes a siloxanyl group, R¹ denotes a substituent with 1 to 20 carbons having polymerizable group, R² to R⁴ respectively and independently denote hydrogen, a substituted or unsubstituted substituent with 1 to 20 carbons, or -X-A, and X denotes a substituted or unsubstituted divalent substituent with 1 to 20 carbons.

6. (Withdrawn) A silicone compound obtained by the process according to Claim 2, wherein the siloxanyl group A is an atomic group represented by the following formula (b),



wherein, A¹ to A¹¹ respectively and independently denote any one of hydrogen, a substituted or unsubstituted alkyl group with 1 to 20 carbon atoms and a substituted or unsubstituted aryl group with 6 to 20 carbons, n denotes an integer of 0 to 200, a, b and c denote respectively and independently an integer of 0 to 20, and n = a = b = c = 0 is not included.


7. (Previously Presented) The process of claim 1, wherein the siloxanyl group A is an atomic group represented by the following formula (b),



wherein, A¹ to A¹¹ respectively and independently denote any one of hydrogen, a substituted or unsubstituted alkyl group with 1 to 20 carbon atoms and a substituted or unsubstituted aryl group with 6 to 20 carbons, n denotes an integer of 0 to 200, a, b and c denote respectively and independently an integer of 0 to 20, and n = a = b = c = 0 is not included.


8. (Previously Presented) The process of claim 7, wherein the siloxanyl group A is selected from the group consisting of tris(trimethylsiloxy)silyl group, bis(trimethylsiloxy)methylsilyl group and trimethylsiloxydimethylsilyl group.

9. (Previously Presented) The process of claim 1, wherein the silicone compound comprises a content of a compound represented by the following general formula (y) in the amount of 0.4% or more and 3% or less,



and a purity of the silicone compound is 87% or more.

10. (Previously Presented) The process of claim 2, wherein the siloxanyl group A is an atomic group represented by the following formula (b),



wherein, A¹ to A¹¹ respectively and independently denote any one of hydrogen, a substituted or unsubstituted alkyl group with 1 to 20 carbon atoms and a substituted or unsubstituted aryl group with 6 to 20 carbons, n denotes an integer of 0 to 200, a, b and c denote respectively and independently an integer of 0 to 20, and n = a = b = c = 0 is not included.